

Fires can benefit plants by disrupting antagonistic interactions

Article (Accepted Version)

García, Y, Castellanos, M C and Pausas, J G (2016) Fires can benefit plants by disrupting antagonistic interactions. *Oecologia*, 182 (4). pp. 1165-1173. ISSN 0029-8549

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/64857/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

[Click here to view linked References](#)

Fires can benefit plants by disrupting antagonistic interactions

Y. García¹· M.C. Castellanos^{1,2}· J.G. Pausas¹

¹ CIDE-CSIC, Montcada, Valencia, Spain

² Current address: School of Life Sciences, University of Sussex, Brighton, BN1 9QG, UK

Corresponding author: Y. García, yedragg@gmail.com, telephone (+34) 96 342 41 62, fax: (+34) 963424160

This study contributes to understand the effects of wildfires on plant-animal interactions considering for the first time different levels of plant-animal specialization on the effects of fire.

Author contribution statement

JGP conceived the idea, JGP and MCC designed the study. MCC and YG collected the data at field. YG performed the statistical analyses and wrote the first version of the manuscript. All authors contributed to the final version of the manuscript.

Abstract

Fire has a key role in the ecology and evolution of many ecosystems, yet its effects on plant-insect interactions are poorly understood. Because interacting species are likely to respond to fire differently, disruptions of the interactions are expected. We hypothesized that plants that regenerate after fire can benefit through the disruption of their antagonistic interactions. We expected stronger effects on interactions with specialist predators than with generalists. We studied two interactions between two Mediterranean plants (*Ulex parviflorus*, *Asphodelus ramosus*) and their specialist seed predators after large wildfires. In *A. ramosus* we also studied the generalist herbivores. We sampled the interactions in burned and adjacent unburned areas during two years by estimating seed predation, number of herbivores and fruit set. To assess the effect of the distance to unburned vegetation we sampled plots at two distance classes from the fire perimeter. Even three years after the fires, *Ulex* plants experienced lower seed damage by specialists in burned sites. The presence of herbivores on *Asphodelus* decreased in burned locations, and the variability in their presence was significantly related to fruit set. Generalist herbivores were unaffected. We show that plants can benefit from fire through the disruption of their antagonistic interactions with specialist seed predators for at least a few years. In environments with a long fire history, this effect might be one additional mechanism underlying the success of fire-adapted plants.

Key-words *Exapion fasciolatum*, generalist, *Horistus orientalis*, seed predation, specialist.

Introduction

Fire is one of the most common disturbances worldwide and can play an important role in the ecology and evolution of many ecosystems (Pausas and Keeley 2009). In environments with a long fire history, such as tropical savannas and Mediterranean ecosystems, fire structures communities and landscapes (Verdú and Pausas 2007; Keeley et al. 2011; Dantas et al. 2013). The effects of fire on plants are relatively well-known (Bond and Van Wilgen 1996; Keeley et al. 2012) and information on the effects of fire on animal populations is also increasing steadily (Swengel 2001; Izhaki 2012; New 2014). Studies on the role of fire in plant-animal interactions have been largely focused on mammal herbivory (e.g., Fuhlendorf et al. 2009; Wan et al. 2014) and, to a lesser extent, on seed predation (e.g., Bond 1984; Andersen 1988; Broncano et al. 2008). However, less is known about how fires disrupt plant-insect interactions and the implications for the plants (Vickery 2002; Knight and Holt 2005; Dafni et al. 2012).

Both antagonistic (e.g. herbivory) and mutualistic (e.g. pollination) interactions between plants and insects are crucial components of natural ecosystems and can determine ecological and evolutionary processes (Herrera and Pellmyr 2002). In ecosystems where wildfires are historically recurrent, many plant species are capable of quickly recovering via resprouting or recruitment from a fire-resistant seedbank (Pausas et al. 2004) and reproduce shortly after the fire. In contrast, fires can directly cause drastic declines in many insect populations, whose recovery then depends on the fire regime and intrinsic characteristics like movement capacity (Swengel 2001; Moretti et al. 2006). Because the different interacting species are likely to respond to fire in varying ways, disruptions of the plant-insect interactions are expected. The dynamics of these disruptions and the postfire recovery of the interactions could therefore have strong consequences for plant populations and constitute important selective pressures for species living in fire-prone environments.

The effects for plants may be different depending on whether fire disrupts mutualistic interactions, potentially decreasing reproductive success, or antagonistic interactions such as herbivory and seed predation. Fires can for example increase seed predation and herbivory when

generalist insects are involved (Andersen 1988; Radho-Toly et al. 2001; Lopes and Vasconcelos 2011). However, there is also evidence of a postfire decrease in insect herbivory in different ecosystems (Whelan and Main 1979; Auld and O'Connell 1989; Vickery 2002; Knight and Holt 2005). In the latter cases, plants could benefit through a release from negative interactions that limit their performance (Hendrix 1988). The reduction of negative effects can be stronger if the interactions involve seed predators, and this can be particularly beneficial for plants in which the success of their first life stages depends on fire.

The consequences of a fire-driven disruption of antagonistic interactions on plant populations will depend, among other factors, on two important interrelated aspects. First, postfire changes in the existing habitat can have a greater impact on species that are tightly dependent on specific habitat characteristics like specialists, compared to generalists (Ewers and Didham 2006). For instance, herbivory and seed predation are often exerted by highly specialized phytophagous insects which interact only with one or a few host plants (Ehrlich and Murphy 1988; Jaenike 1990). Consequently, the alteration of their host plants may lead to changes in their abundance and distribution (Larsson et al. 2000). The recolonization of burnt areas by generalist animals might therefore be faster than by specialists, because the specialist's mobility into the interior of the burnt may be restricted to the presence and the regeneration of their only host. Second, the distance to unburned vegetation might also have an impact on the speed of recolonization and therefore on the duration of the disruption, which can result in spatial variation in the interaction from the edge towards the interior of the burned area. The contrast between specialists and generalists may be even stronger in large fires where species have to migrate long distances to reach the center of the burned area. A stronger disruption of specialized interactions compared to generalist ones would thus be expected, and this effect could be exacerbated as one moves from the edge to the center of the fire. While some previous studies show that fire may modify the plant-insect interaction, less is known about the implications for the plant's reproductive performance (e.g., Whelan and Main 1979; Auld and O'Connell 1989; Vickery 2002), and particularly on how these implications differ depending on the level of specialization of the insect.

Our hypothesis is that plants that quickly regenerate after fire may additionally benefit from it because fire disrupts antagonistic interactions, and that this effect will be exacerbated with the distance to the unburned vegetation. To test it, we studied two interactions between plants and their specialist seed predators after recent wildfires in Mediterranean shrublands of eastern Spain: (1) the Mediterranean gorse *Ulex parviflorus* and its seed predator, the weevil *Exapion fasciolatum*; and (2) the branched asphodel *Asphodelus ramosus* and the specialist mirid bug *Horistus orientalis*, together with other generalist insects. In the two systems the life cycles of the specialist insects depend entirely on their host plants. We expected that fires would have strong negative effects on the local populations of both specialist predators, and would in turn benefit the host plants by reducing seed predation.

Material and methods

Plant-insect interaction I: *Ulex parviflorus*-*Exapion fasciolatum*

The Mediterranean gorse, *Ulex parviflorus* Pourr. (Fabaceae) is a spiny perennial shrub from the western Mediterranean Basin. It can live up to 25 years (Baeza and Vallejo 2006) and, as observed in this study, individuals can reach their mature stage as soon as two years after fire. One or two (occasionally more) seeds develop inside small pods and are dispersed explosively at the beginning of the summer. A preliminary analysis suggested that the variance in the number of seeds per pod is not related to contrasted fire regimes (mean number of seeds/pod = 1.29 and 1.34 in populations growing under high or low fire frequency respectively, N= 3206 pods examined). *Ulex parviflorus* is common in fire-prone Mediterranean shrublands where it recruits massively after fire, when high soil temperatures break seed dormancy and induce germination (postfire obligate seeder; Paula et al. 2009; Moreira et al. 2010; Moreira and Pausas 2012).

Ulex parviflorus seeds are attacked by the weevil *Exapion fasciolatum* Wagner (Brentidae: Apioninae). Information on this species is scarce, but *Exapion* species are specialist predators of the Genisteae tribe (Fabaceae; Alonso-Zarazaga 1990), including *Ulex* (Barat et al. 2007; Tarayre et al. 2007). In the closely related *Exapion ulicis*-*Ulex europaeus* interaction, weevil

predation can damage up to 90% of the gorse pods and may explain phenological shifts in the plant to reduce predation impacts (Barat et al. 2007; Tarayre et al. 2007). The weevil's life cycle depends entirely on the host plant. In early spring, females lay their eggs inside the gorse ovaries or small green pods, where larvae and pupa develop while feeding on the seeds. Adults emerge with ripe pod dehiscence. *Ulex parviflorus* pods can also contain a parasitoid wasp (*Eurytoma* sp.) feeding on the larvae and pupae of *Exapion fasciolatum*.

Plant-insect interaction II: *Asphodelus ramosus*-*Horistus orientalis*

Asphodelus ramosus L. (= *A. aestivus* Brot., Liliaceae) is a Mediterranean geophyte widely distributed along the Mediterranean basin (Lifante 1996). It has a short rhizome surrounded by tubers and a basal rosette of leaves that produces a branched flowering scape. Thanks to the resprouting capacity from the rhizome, this species is favored by heavy grazing and recurrent fires (Pantis and Margaritis 1988); in fact, it flowers massively in burned areas.

Horistus orientalis Gmelin (= *Capsodes lineolatus* Br., Hemiptera: Miridae) is a phytophagous bug that feeds on *Asphodelus ramosus*. Members of the mirid family exhibit a high degree of host-plant specificity (Cassis and Schuh 2012) and in our study region this species has never been confirmed feeding on other plants (Luis Vivas pers. comm). Mirids often develop synchronously with the plant, from the deposition of the eggs within the scape tissues, until adult emergence after the nymph stage (Wheeler 2001; Cassis and Schuh 2012). Published information about *H. orientalis* is very limited, but our observations suggest that its entire life cycle occurs on the plant, as in the closely related bug *Capsodes infuscatus*. The eggs are deposited inside the inflorescence stalk in the spring and adults disperse the next spring (Ayal and Izhaki 1993; Izhaki et al. 1996). Both nymphs and adults feed on leaves and especially on flowers and fruits. In the case of *C. infuscatus*, the damage produced on *A. ramosus* can reach 100% of fruit loss (Ayal and Izhaki 1993). *Asphodelus ramosus* is also attacked by generalist herbivores; we mainly observed two phytophagous beetles from the subfamily Cetoniinae, *Tropinota squalida* Scop. and *Oxythyrea funesta* Poda, and the bug *Carpocoris fuscipinus* Boh. (Pentatomidae), among others.

Study sites and sampling

Each interaction was studied during two consecutive years in two burned locations from different wildfires in Eastern Spain (Valencia; see Table 1). The region shows a typical Mediterranean climate with frequent fires (Pausas 2004; Pausas and Paula 2012). To study the effect of fire on the interactions we sampled plots inside each burned location plus unburned adjacent (control) plots where no fires have been registered for at least 20 years. Unburned plots were carefully chosen to be representative of the pre-fire conditions (e.g. same plant species composition, dominant species and soil characteristics), and when possible, close to fire perimeter. The same control and burned plots were sampled during the two years of the study when possible. To assess the effect of the distance to unburned vegetation on the interactions, plots were assigned to three different categories according to their distance to the fire's perimeter: (a) control plots in the adjacent unburned areas ("Unburned"), (b) plots located inside the burned area and up to 500 meters (mean= 268) from the fire perimeter ("Edge") and (c) plots at more than 500 meters (mean= 1199, maximum= 2400) from the fire perimeter ("Center"). Plots at burned areas were carefully selected to avoid the proximity of unburned patches. Distances were estimated using geographic information tools and digital maps provided by the regional government of Valencia.

Seed predation on *Ulex parviflorus* was measured in 48 plots in two locations where large wildfires (of more than 20000 ha each) had occurred in the summer of 2012: Cortes de Pallás (hereafter, Cortes) and Andilla, both in the province of Valencia (Table1). Before the fires, all plots were Mediterranean shrublands dominated by *Rosmarinus officinalis*, *Ulex parviflorus*, several *Cistus* species and *Quercus coccifera*. Field work was carried out between late March and June in 2014 and 2015 and corresponded to the first two postfire flowering years for the newly recruited individuals of *U. parviflorus*. The sampling included 34-35 burned plots and 13-14 plots in the adjacent unburned areas (see Table S1 in Supplementary Material for details on plots at each fire location). At each plot, 400 mature pods were collected from 10 haphazardly chosen *U. parviflorus* plants (40 pods per plant), separated from each other by at

least 5 m. The 400 pods from Andilla's burned plots in 2014 were collected from 20 plants (20 per plant) because fruit production per plant was lower in that location. We chose a fixed number of pods per plant to estimate the levels of seed predation as opposed to attempting to quantify whole-plant production, a difficult task given the massive and extended flowering period of *U. parviflorus*. Overall the total number of pods sampled was 18370 in 2014 and 19265 in 2015.

The content of each pod was observed at the laboratory under a stereo-microscope. The presence of the weevil at larva, pupa or adult stage was recorded in each pod. When a parasitoid wasp was observed inside the pod, it was also counted as predated, i.e., we assumed that wasps had emerged from a weevil larva (Barat et al. 2007). We used the proportion of predated pods as a measure of the effect of the seed predator on the fitness of the plant. This method directly estimates weevil predation within each pod and allows to differentiate their effect from other predispersal predators as mentioned above (Barat et al. 2007).

The study on *A. ramosus* was conducted in Cortes and in a second smaller wildfire that occurred in February 2014 in Segorbe (province of Castellón; Table1). Sampling was performed during spring when *A. ramosus* was already in bloom, and included a total of 15 plots in 2014 (9 burned and 6 unburned plots) and 14 in 2015 (8 burned and 6 unburned plots, for details see Table S2 in Supplementary Material). At each plot, the presence and activity of the specialist bug *Horistus orientalis* and the most abundant herbivores (Cetoniinae and Pentatomidae) were recorded on 50 haphazardly chosen *Asphodelus* plants separated from each other by at least 5 m; censuses were conducted between 10:00 and 16:00 h. Other generalist herbivores were observed only rarely and were thus not included in the analyses. The number of branches, floral buds and flowers were also recorded for each plant. At the end of the flowering season (May-June) we collected ripening fruits from all plants and counted healthy seeds in the laboratory in all plots within the burned areas. The proportion of fruits in relation to the number of flowers produced (fruit set) was considered an indicator of reproductive success and was analyzed with respect to the presence of the seed predators on the plant (see below). We used fruit set as a proxy for reproductive success because it corrects for variation in plant size as opposed to using the

absolute number of fruits or seeds produced per plant. Although fruit set is not a direct measure of the plant damage, it may reflect the total effects of the different feeding habits of the insects including green parts of the plant and also fruits. In fact there is evidence of a strong negative relationship between plant fruit set and the bug's abundance in the closely related *Asphodelus aestivus*-*Capsodes infuscatus* interaction (Ayal and Izhaki 1993; Izhaki et al. 1996). Fires could also affect other factors linked to plant fruit set such as resource availability and pollination. We expect a limited effect of pollination, because this species has a generalized pollination system (Lifante 1996; Lázaro et al. 2016) and flying pollinators tend to recover quickly after fires (Potts et al. 2003). The increase in resources often associated with postfire environments could also have positive effects on fruit production. However, we did not find a difference either in the number of flowers nor in the absolute seed production between burned and unburned sites (see results), suggesting a limited relevance of the potential changes in resources.

Statistical analysis

To examine whether seed predation on *U. parviflorus* at the plant level differed between burned and unburned plots, we used a generalized linear mixed model (GLMM), with a binomial error distribution. For each year of sampling, the GLMM included the burned vs. unburned treatment as a fixed factor and plot nested within locality as random factor. The same approach was used to test whether seed predation varied between Edge and Center zones (i.e., within the burned area).

To test whether the number of specialist *Horistus orientalis* individuals differed between *A. ramosus* plants from burned and control plots, we used a similar GLMM model structure as above, in this case with a Poisson error distribution. We also used the same model structure to test for differences in the number of generalist herbivores (Pentatomidae plus Cetoninae), in the total herbivores (*H. orientalis* and generalists together) and in the number of flowers and seeds produced per plant. We then tested for distance effects in the number of the three insect groups (specialist, generalist and total) by fitting a GLMM to the distance class variable (Edge vs Center). To evaluate to what extent the variability observed in herbivores in the burned zone

correlates with the variability in plant fitness, we fitted the fruit set against the number of *Horistus* bugs using a GLMM with a binomial error distribution. For this, we used the number of *Horistus* bugs in relation to the number of flowers of each plant, and tested it with the nested design mentioned above to account for plot variability. We used a similar model with the number of total herbivores (also corrected by the number of flowers). For both studied interactions we also fitted a GLMM that included the combined data of both sampling years and the year as a random factor. Overdispersion was tested and corrected when necessary. All models were run with 'lme4' package in R (Bates et al. 2014).

Results

Ulex plants from burned plots showed a much lower proportion of predated pods by their specialist seed predator (<5%) than the adjacent unburned plots (>15%; Table 2, Fig. 1 and Table S1 in Supplementary Material). The number of *Exapion* weevils was also lower in burned plots and, unexpectedly, decreased on the second sampling year (Table 2 and Table S1). The predation of *Ulex* pods and the number of weevils decreased from the Edge to the Center of the burnt; this decrease was significant for 2014 and for the overall period, but not for 2015 (Table 2, Fig. 1 and Table S1 in Supplementary Material).

The number of specialist *H. orientalis* bugs as well the total number of herbivores on *Asphodelus* plants were significantly lower in burned than in the unburned plots on both years of sampling (Table 2, Fig. 2 and Fig. S1 in Supplementary Material). However, the number of generalist herbivores did not differ neither between the two environments nor between the Edge and the Center (Table 2, Fig. 2). Neither the number of flowers nor seeds produced per plant showed significant differences between unburned and burned plots ($N=1414$, $P=0.09$ and $P=0.54$ respectively). *Asphodelus* fruit set showed a significant negative relationship with the abundance of *Horistus* and also with the abundance of total herbivores when data from the two years were combined (see Fig. 3).

Discussion

Previous studies on the responses of insect herbivory to fire have shown an increment in herbivory related to a post-fire increase in herbivore abundance (Andersen 1988; Radho-Toly et al. 2001; Lopes and Vasconcelos 2011). Typically these studies have focused on generalized interactions where the insects do not depend on the recovery of a specific plant to recolonize the burned areas. Our study, however, shows that fires can instead benefit plants by decreasing their herbivory pressure particularly from specialist insects (Auld and O'Connell 1989, Vickery 2002). Even three years after the fire, *Ulex parviflorus* plants experienced lower seed damage in burned plots than in control unburned sites (Table 2, Fig. 1). For *Asphodelus ramosus*, we found that the variability in fruit set in burned areas is significantly related to the presence of the specialist bug. That is, the disruption of the specialized interaction can affect plant fitness. In plant species with a quick recovery after fires, this disruption also coincides with an increase of resources and reduced competition after fire. Thus, the evidence suggests that the disruption of antagonistic interactions between plants and insects following a fire might be one mechanism contributing to plant success in fire-prone ecosystems.

Many insect populations decline immediately after a fire; furthermore, fire temporarily decreases the presence of the host plant required for insect development. Both the drop in insect populations and the reduction in food resources may have a stronger impact on specialist than on generalist insects (Swengel 1996, 1998, 2001). Generalist herbivores have access to a wider range of recovering plants, thus showing a faster re-colonization than the specialists. Consistently, in this study the presence of two specialist predators was much lower in plants from burned plots while generalist herbivores recorded on *A. ramosus* remained unaffected.

After three years of the wildfires we did not find a consistent effect of the distance from the perimeter of the fire on seed predation, herbivory pressures or plant fitness (Table 2). Several previous studies had shown a reduction of herbivory with the distance to unburned vegetation. For example, *Banksia* and *Eucalyptus* seedlings experienced a lower damage by generalist grasshoppers in large burned areas compared with small ones (Whelan and Main 1979). In a sandhill ecosystem, plants from the center of a burnt suffered half of the impact of insect herbivory compared to plants from the fire's edge (Knight and Holt 2005). The limited distance

effect in the present study may be explained by the low postfire predation levels on *U. parviflorus* and the low number of *H. orientalis* individuals recorded in most burned plots. In fact, we did detect a significant decrease in *Ulex* predation towards the center of the burned zone in 2014 (Table 2), when the predation by the weevil was 5 times higher than in 2015. The causes behind the decreased densities in specialist predators are unknown, and seem to be unrelated to climate conditions, which did not show major differences between the two sampling years. Despite there is little knowledge on the population dynamics of the studied specialist insects, our results suggest that fires may have a great negative effect on their populations and it may last for several years while recolonization takes place.

The disruption of the interaction by fire is likely to have long-term benefits for the plant. *Ulex parviflorus* benefits from fire because the heat reached during a fire breaks seed dormancy and greatly stimulates germination from the soil seedbank (Paula et al. 2009; Moreira et al. 2010; Moreira and Pausas 2012), and thus the postfire population size is greater than in prefire populations. *Asphodelus* can also take advantage of the fire because the canopy gap opened allows this species to flower profusely, otherwise the high density of the shrubland limits sexual reproduction (Pantis and Margaris 1988; Pantis and Mardiris 1992). Here we show that fires can generate an additional benefit to the plant by creating a window of opportunity for reproduction under a lower predation pressure from their specialist herbivores. Although this release is likely temporary, it may have long-term effects because it ensures the quick refill of the seedbank after fire and thus the ability to massively recruit even under short fire intervals. For *Asphodelus*, it increases fruit set and promotes sexual reproduction under suitable postfire recruitment conditions. To what extent these beneficial effects can be generalized to other fire-adapted plant species reminds to be studied; previous research on this regard is limited to single populations or to prescribed fire regimes (Auld and O'Connell 1989; Vickery 2002).

Additionally to predispersal predation, fires can affect postdispersal seed predation which may also have implications on plant fitness (Andersen 1988; Ordoñez Retana 2004; Zwolak et al. 2010; Keeley et al. 2012). For example, seed predation on *Pinus* species from the Mediterranean increased after fires coinciding with a high presence of ants and rodents

(Ordoñez and Retana 2004; Broncano et al. 2008). These changes on secondary seed predation are related with the dispersal season and the time since fire (Ordoñez and Retana 2004). Our two studied plants disperse seeds from spring (*Ulex parviflorus*) to summer (*Asphodelus ramosus*), when ants can be especially abundant after fires (Ordoñez and Retana 2004). Although ants are known as the main seed predators in burned zones (Rey 2002; Broncano et al. 2008), there is evidence that *Ulex parviflorus*, which has elaiosomes, can show higher germination rates after ant-dispersal (López-Vila and García-Fayos 2005). We are not aware of any information on seed predation by ants in *Asphodelus ramosus*. In any case, further studies depicting the relative role of ants as predators and dispersers (Auld and Denham 1999) on the two studied plants at burned areas would contribute to better understanding their success in burning ecosystems.

Despite the accepted key role of fire in many ecosystems, the responses of plant-insect interactions to fire are not well known (Dafni et al. 2012). This gap in the knowledge is even more remarkable for fire-prone ecosystems such as the Mediterranean ones, where there is evidence of fire-adaptive traits in many different species (Keeley et al. 2011). Our study, for which we monitored two plant-insect interactions across several years at different locations, indicates that when fire has disrupted specialized antagonistic interactions between insects and plants capable of quickly regenerating after fire, these plants can benefit from this for several years. This “cleaning” effect by fire might be one of the factors promoting the success of fire-adapted plants. It remains to be determined how wide-spread these effects are across different ecological settings. However, we feel that the fact that we observed beneficial effects for two plant species - interacting with different insects after multiple fires at two different locations in two consecutive years - suggests that it may be a general phenomenon. However, fires will not always benefit plants as they can also disrupt mutualisms (Dafni et al. 2012) and change the dispersal-predation balance of generalized interactions (Andersen 1988; Radho-Toly et al. 2001; Ordoñez and Retana 2004; Broncano et al. 2008; Lopes and Vasconcelos 2011). The current crisis of biotic interactions and the expected increase in fire size and frequency associated with

anthropogenic activities, make understanding the effects of fire on plant-insect interactions an urgent need.

Acknowledgements

We thank M.A. Alonso-Zarazaga, J. Baixeras, J.V. Falcó-Gari, L. Vivas, S.M. Alario and M.A. Orrico for insect identification. We are grateful to Abel Rubira, Santiago Donat-Caerols, Miguel Adán, Alejandro (Dani) Ginart, Beatriz López Gurillo, Lucía Tortajada and José Vicente Pérez for their help in the field and in the laboratory. CIDE is a joint institute of the Spanish National Research Council (CSIC), the University of Valencia and the regional government of Valencia (*Generalitat Valenciana*). *Generalitat Valenciana* also provided detailed fire maps. This research was financed by projects TREVOL and FILAS (CGL2012-39938-C02-01, CGL2015-64086) from the Spanish Government and PROMETEO/2016/021 from *Generalitat Valenciana*. Y.G. was supported by an FPI scholarship (BES-2013-062728) from the Spanish Ministry of Economy and Competitiveness.

Conflict of interest

We declare no conflict of interest

References

- Alonso-Zarazaga MA (1990) Revision of the supraspecific taxa in the Palaearctic Apionidae Schoenherr, 1823 (Coleoptera, Curculionoidea). 2. Subfamily Apioninae Schoenherr, 1823: introduction, keys and descriptions. *Graellsia* 46:19–156.
- Andersen AN (1988) Immediate and longer-term effects of fire on seed predation by ants in sclerophyllous vegetation in south-eastern Australia. *Aust J Ecol* 13:285–293. doi:10.1111/j.1442-9993.1988.tb00976.x
- Auld TD, Denham AJ (1999) The role of ants and mammals in dispersal and post-dispersal seed predation of the shrubs *Grevillea* (Proteaceae). *Plant Ecol* 144:201–213. doi:10.1023/A:1009817132378
- Auld TD, O'Connell MA (1989) Changes in predispersal seed predation levels after fire for two Australian legumes, *Acacia elongata* and *Sphaerolobium vimineum*. *Oikos* 54:55–59. doi:10.2307/3565896
- Ayal Y, Izhaki I (1993) The effect of the mirid bug *Capsodes infuscatus* on fruit production of the geophyte *Asphodelus ramosus* in a desert habitat. *Oecologia* 93:518–523. doi:10.1007/BF00328960
- Baeza MJ, Vallejo VR (2006) Ecological mechanisms involved in dormancy breakage in *Ulex parviflorus* seeds. *Plant Ecol* 183:191–205. doi:10.1007/s11258-005-9016-0
- Barat M, Tarayre M, Atlan A (2007) Plant phenology and seed predation: interactions between gorses and weevils in Brittany (France). *Entomol Exp Appl* 124:167–176. doi:10.1111/j.1570-7458.2007.00565.x
- Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1-7, <http://CRAN.R-project.org/package=lme4>.
- Bond WJ (1984) Fire survival of Cape Proteaceae-influence of fire season and seed predators. *Plant Ecol* 56: 65–74. doi:10.1007/BF00033048
- Bond WJ, van Wilgen BW (1996) *Fire and Plants*, 1st edn. Chapman and Hall, London.

394 Broncano MJ, Rodrigo A, Retana J (2008) Post-dispersal seed predation in *Pinus halepensis* and
 395 consequences on seedling establishment after fire. *Int J Wildland Fire* 17:407–414.
 396 doi:0.1071/WF07095

397 Cassis G, Schuh RT (2012) Systematics, biodiversity, biogeography, and host associations of the
 398 Miridae (Insecta: Hemiptera: Heteroptera: Cimicomorpha). *Annu Rev Entomol* 57:377–
 399 404. doi:10.1146/annurev-ento-121510-133533

400 Dafni A, Izhaki I, Ne'eman G (2012) The effect of fire on biotic interactions in Mediterranean
 401 basin ecosystems: pollination and seed dispersal. *Isr J Ecol Evol* 58:235–250.
 402 doi:10.1560/IJEE.58.2-3.235

403 Dantas VL, Batalha MA, Pausas JG (2013) Fire drives functional thresholds on the savanna-
 404 forest transition. *Ecology* 94:2454–2463. doi:10.1890/12-1629.1

405 Ehrlich PR, Murphy DD (1988) Plant chemistry and host range in insect herbivores. *Ecology*
 406 69:908–909. doi:10.2307/1941244

407 Ewers RM, Didham RK (2006) Confounding factors in the detection of species responses to
 408 habitat fragmentation. *Biol Rev Camb Philos Soc* 81:117–142.
 409 doi:10.1017/S1464793105006949

410 Fuhlendorf SD, Engle DM, Kerby JAY, Hamilton R (2009) Pyric herbivory: rewilding
 411 landscapes through the recoupling of fire and grazing. *Conserv Biol* 23:588–598.
 412 doi:10.1111/j.1523-1739.2008.01139.x

413 Hendrix SD (1988) Herbivory and its impact on plant reproduction. In: Lovett Doust J, Lovett
 414 Doust L (eds) *Plant reproductive ecology*. Oxford University Press, Oxford, pp 246–
 415 263.

416 Herrera CM, Pellmyr O (2002) *Plant animal interactions: an evolutionary approach*, 1st edn.
 417 Blackwell Science, Oxford.

418 Izhaki I (2012) The impact of fire on vertebrates in the Mediterranean Basin: an overview. *Isr J*
 419 *Ecol Evol* 58:221–233. doi:10.1560/IJEE.58.2-3.221

420 Izhaki I, Maestro N, Meir D, Broza M (1996) Impact of the Mirid bug *Capsodes infuscatus*
 421 (Hemiptera: Miridae) on fruit production of the geophyte *Asphodelus aestivus*: the
 422 effect of plant density. Fla Entomol 79:510–520. doi:10.2307/3496063

423 Jaenike J (1990) Host specialization in phytophagous insects. Annu Rev Ecol Evol Syst 21:243–
 424 273. doi:10.1146/annurev.es.21.110190.001331

425 Keeley JE, Bond WJ, Bradstock RA, Pausas JG, Rundell PW (2012) Fire in Mediterranean
 426 ecosystems: ecology, evolution and management, 1st edn. Cambridge University Press,
 427 Cambridge.

428 Keeley JE, Pausas JG, Rundel PW, Bond WJ, Bradstock RA (2011) Fire as an evolutionary
 429 pressure shaping plant traits. Trends Plant Sci 16:406–411.
 430 doi:10.1016/j.tplants.2011.04.002

431 Knight TM, Holt RD (2005) Fire generates spatial gradients in herbivory: an example from a
 432 Florida sandhill ecosystem. Ecology 86:587–593. doi:10.1890/04-1069

433 Larsson S, Ekbohm B, Björkman C (2000) Influence of plant quality on pine sawfly population
 434 dynamics. Oikos 89:440–450. doi:10.1034/j.1600-0706.2000.890303.x

435 Lázaro A, Tscheulin T, Devalez J, Nakas G, Petanidou T (2016) Effects of grazing intensity on
 436 pollinator abundance and diversity, and on pollination services. Ecol Entomol. In press.
 437 doi:10.1111/een.12310

438 Lifante ZD (1996) Reproductive biology of *Asphodelus aestivus* (Asphodelaceae). Plant Syst
 439 Evol 200:177–191. doi:10.1007/BF00984934

440 Lopes CT, Vasconcelos HL (2011) Fire increases insect herbivory in a neotropical savanna.
 441 Biotropica 43:612–618. doi:10.1111/j.1744-7429.2011.00757.x

442 López-Vila JR, García-Fayos P (2005) Diplochory in *Ulex parviflorus* Pourr. Acta Oecol
 443 28:157–162. doi:10.1016/j.actao.2005.03.008

444 Moreira B, Pausas JG (2012) Tanned or burned: the role of fire in shaping physical seed
 445 dormancy. PLoS ONE 7:e51523. doi:10.1371/journal.pone.0051523

- Moreira B, Tormo J, Estrelles E, Pausas JG (2010) Disentangling the role of heat and smoke as germination cues in Mediterranean Basin flora. *Ann Bot* 105:627–635. doi:10.1093/aob/mcq017
- Moretti M, Duelli P, Obrist, MK (2006) Biodiversity and resilience of arthropod communities after fire disturbance in temperate forests. *Oecologia* 49:312–327. doi:10.1007/s00442-006-0450-z
- New TR (2014) *Insects, fire and conservation*, 1st edn. Springer, New York.
- Ordóñez JL, Retana J (2004) Early reduction of post fire recruitment of *Pinus nigra* by post dispersal seed predation in different time since fire habitats. *Ecography* 27:449–458. doi:10.1111/j.0906-7590.2004.03886.x
- Pantis JD, Mardiris TA (1992) The effects of grazing and fire on degradation processes of Mediterranean ecosystems. *Israel J Bot* 41:233–242. doi:10.1080/0021213X.1992.10677230
- Pantis JD, Margaris NS (1988) Can systems dominated by asphodels be considered as semi-deserts? *Int J Biometeorol* 32:87–91. doi:10.1007/BF01044899
- Paula S, Arianoutsou M, Kazanis D, Tavsanoğlu Ç, Lloret F, Buhk C, Ojeda F, Luna B, Moreno JM, Rodrigo A, Espelta JM, Palacio S, Fernández-Santos B, Fernandes PM, Pausas JG (2009) Fire-related traits for plant species of the Mediterranean Basin: Ecological Archives E090-094. *Ecology* 90:1420–1420. doi:0.1890/08-1309.1
- Pausas JG (2004) Changes in fire and climate in the eastern Iberian Peninsula (Mediterranean basin). *Climatic Change* 63:337–350. doi:10.1023/B:CLIM.0000018508.94901.9c
- Pausas JG, Bradstock RA, Keith DA, Keeley JE (2004) Plant functional traits in relation to fire in crown-fire ecosystems. *Ecology* 85:1085–1100. doi:10.1890/02-4094
- Pausas JG, Keeley JE (2009) A burning story: the role of fire in the history of life. *BioScience* 59:593–601. doi:10.1525/bio.2009.59.7.10
- Pausas JG, Paula S (2012) Fuel shapes the fire-climate relationship: evidence from Mediterranean ecosystems. *Global Ecol Biogeogr* 21:1074–1082. doi:10.1111/j.1466-8238.2012.00769.x

- Potts SG, Vulliamy B, Dafni A, Ne'eman G, O'Toole C, Roberts S, Willmer P (2003) Response of plant-pollinator communities to fire: changes in diversity, abundance and floral reward structure. *Oikos* 101:103–112. doi:10.1034/j.1600-0706.2003.12186.x
- Radho Toly S, Majer JD, Yates C (2001) Impact of fire on leaf nutrients, arthropod fauna and herbivory of native and exotic eucalypts in Kings Park, Perth, Western Australia. *Austral Ecol* 26:500–506. doi:10.1046/j.1442-9993.2001.01133.x
- Rey PJ, Garrido JL, Alcántara JM, Ramírez JM, Aguilera A, García L, Manzaneda AJ, Fernández R (2002) Spatial variation in ant and rodent post-dispersal predation of vertebrate-dispersed seeds. *Funct Ecol* 16:773–781. doi:10.1046/J.1365-2435.2002.00680.X
- Swengel AB (1996) Effects of fire and hay management on abundance of prairie butterflies. *Biol Conserv* 76:73–85. doi:10.1016/0006-3207(95)00085-2
- Swengel AB (1998) Effects of management on butterfly abundance in tallgrass prairie and pine barrens. *Biol Conserv* 83:77–89. doi:10.1016/S0006-3207(96)00129-2
- Swengel AB (2001) A literature review of insect responses to fire, compared to other conservation managements of open habitat. *Biodivers Conserv* 10:1141–1169. doi:10.1023/A:1016683807033
- Tarayre M, Bowman G, Schermann-Legionnet A, Barat M, Atlan A (2007) Flowering phenology of *Ulex europaeus*: ecological consequences of variation within and among populations. *Evol Ecol* 21:395–409. doi:10.1007/s10682-006-9109-9
- Verdú M, Pausas JG (2007) Fire drives phylogenetic clustering in Mediterranean Basin woody plant communities. *J Ecol* 95:1316–1323. doi:10.1111/j.1365-2745.2007.01300.x
- Vickery PD (2002) Effects of the size of prescribed fire on insect predation of northern blazing star, a rare grassland perennial. *Conserv Biol* 16:413–421. doi:10.1046/j.1523-1739.2002.00494.x
- Wan HY, Rhodes AC, St Clair SB (2014) Fire severity alters plant regeneration patterns and defense against herbivores in mixed aspen forests. *Oikos* 12:1479–1488. doi:10.1111/oik.01521

502 Wheeler AG (2001) Biology of the plant bugs (Hemiptera: Miridae): pests, predators,
503 opportunists, 1st edn. Cornell University Press, New York.
504 Whelan RJ, Main AR (1979) Insect grazing and post fire plant succession in south west
505 Australian woodland. Aust J Ecol 4:387–398. doi:10.1111/j.1442-9993.1979.tb01566.x
506 Zwolak R, Pearson DE, Ortega YK, Crone EE (2010) Fire and mice: Seed predation moderates
507 fire's influence on conifer recruitment. Ecology 91:1124–1131. doi:10.1890/09-0332.1

508

509 **Electronic Supplementary Material**

510 **Table S1.** Number of predated and total pods collected of *Ulex parviflorus*, total number of
511 *Exapion fasciolatum* individuals per plot, and the distance class for each plot.

512

513 **Table S2.** Number of specialist *Horistus orientalis* bugs, generalists and total herbivores
514 recorded on *Asphodelus ramosus*, mean fruit set and total number of seeds at each plot.

515

516 **Figure S1.** Number of total herbivores on *Asphodelus ramosus* plants from unburned and
517 burned plots in two years.

518 **Table 1.** Fire location name, year of fire occurrence and sampling years for each study system:
 519 a) *Ulex parviflorus* and the specialist weevil *Exapion fasciolatum* and b) *Asphodelus ramosus*
 520 and the specialist bug *Horistus orientalis* and its generalist herbivores.

System	Fire location	Year	Sampling years
<i>U. parviflorus-E. fasciolatum</i>	Cortes	2012	2014 & 2015
	Andilla	2012	2014 & 2015
<i>A. ramosus-herbivores</i>	Cortes	2012	2014 & 2015
	Segorbe	2014	2014 & 2015

Table 2. Results from generalized linear mixed models of the effects of fire on seed predation and herbivory in two Mediterranean plants (*Ulex parviflorus* and *Asphodelus ramosus*). For each response variable and year of study, we first compared plants from Unburned vs Burned (U. vs B.) plots, and then for the burned plots, we compared Edge vs Center plots (E. vs C.). For *U. parviflorus*, the models test for differences in the incidence of seed predation by the weevil *Exapion fasciolatum* and the number of *E. fasciolatum* weevils. For *A. ramosus*, response variables were the number of *Horistus orientalis* (specialist bug), the number of generalist herbivores, and the total number of herbivores. The table shows the sample sizes (N) and, for each significant model, the estimated parameter for the fixed effects (Estimate) and the associated significance (*P*, n.s, not significant; **P* < 0.05, ***P* < 0.01, ****P* < 0.001). Estimate refers to the coefficient of Burned (in relation to Unburned) and of the Center (in relation to the Edge).

		2014			2015			2014 & 2015		
	Test	N	Estimate	<i>P</i>	N	Estimate	<i>P</i>	N	Estimate	<i>P</i>
<i>Ulex parviflorus</i>										
Predated pods	U. vs B.	578	-1.46	***	469	-1.73	***	1047	-1.60	***
	E. vs C.	480	-0.56	**	334	-	n.s	814	-0.45	*
Specialist weevil	U. vs B.	578	-1.38	***	469	-1.42	***	1047	-1.40	***
	E. vs C.	480	-0.47	**	334	-	n.s	814	-0.45	**
<i>Asphodelus ramosus</i>										
Specialist bug	U. vs B.	729	-1.90	***	686	-1.60	**	1415	-1.77	***
	E. vs C.	433	-	n.s	400	-	n.s	833	-	n.s
Generalist herbivores	U. vs B.	729	-	n.s	686	-	n.s	1415	-	n.s
	E. vs C.	433	-	n.s	400	-	n.s	833	-	n.s
Total herbivores	U. vs B.	729	-0.38	*	686	-1.19	**	1415	-0.61	**
	E. vs C.	433	-	n.s	400	-	n.s	833	-	n.s

Figure legends

Fig. 1. Predation rate of *Ulex parviflorus* pods in unburned plots (grey box) and burned plots (white boxes, “Edge” and “Center”) for two years of sampling. N= 13 and 14 “Unburned” plots, and N= 15 and 14 “Edge” plots in 2014 and 2015 respectively. N= 20 for “Center” plots on both years.

Fig. 2. Number of specialist bugs *Horistus orientalis* on *Asphodelus ramosus* plants from unburned (grey box) and burned plots (white boxes in categories “Edge” and “Center”) in two years. The number of individuals was estimated in 50 *A. ramosus* plants per plot (N= 6 unburned and N= 4 “Center” plots in the two studied years. N= 5 and 4 “Edge” plots in 2014 and 2015).

Fig. 3. *Asphodelus ramosus* fruit set (proportion of fruits in relation to flowers) in relation to the number of the specialist bug *Horistus orientalis* in plants from burned plots. The relation is significant either considering *Horistus* only (N= 833, Estimate= -16.47, $P= 0.018$) or total herbivores (N= 833, Estimate= -12.57, $P= 0.018$) in a GLMM considering random effects and correcting for plant size.

Proportion of predated pods per plant




Fig.2





Fig.3





[Click here to access/download](#)

Supplementary Material

Supplementary_material_replay.pdf

